arXiv:1612.06500v2 [math.DG] 5 Jan 2017

Minimal submanifolds in certain types of kaehler product manifold

Xingda Liu Bang Xiao *

Department of Mathematics, Chongqing University of Technology,

Chongqing 400054, The People's Republic of China

(xingdaliu0924@cqut.edu.cn bangxiao@cqut.edu.cn)

January 6, 2017

Abstract

Let M be a real l-dimensional minimal submanifold with flat normal connection in a kaehler product manifold $\overline{M}^m \times \overline{M}^n$ where $\overline{M}^m$ and $\overline{M}^n$ are complex m-dimensional and complex n-dimensional kaehler manifolds with constant holomorphic sectional curvature c_1 and c_2 respectively. We give a formula for the Laplacian of the second fundamental form of M. Specially we discuss the F-anti invariant case. We also give some applications of this formula.

Key words: Kaehler product manifold, minimal submanifolds.

2000 Mathematics Subject Classification: 58A05; 53C55.

1 Introduction

The complex projective space CP^m and its submanifolds have been studied by many researchers. For the product manifold K.Yano and M.Kon [7] give the general results. The F-invariant submanifolds, invariant submanifolds, totally real submanifolds, of Kaehler product manifold have been studied, for example [1], [2], [6], [3].

In this paper we give a formula for the Laplacian of the second fundamentai form of a l-dimensional minimal submanifold with flat normal connection in a kaehler product manifold $\overline{M}^m \times \overline{M}^n$, where $\overline{M}^m$ and $\overline{M}^n$ are complex m-dimensional and complex n-dimensional kaehler manifolds with constant holomorphic sectional curvature c_1 and c_2 respectively. We also give some applications of this formula. Specially in the F-anti invariant case, we get several simpler results.

*Corresponding author.Supported by the NNSF 11526040 of China. Partially supported by the NNSF 11371386 of China

2 Preliminaries

Let $\overline{M}^m$ be a Kaehlerian manifold of complex dimension m with constant holomorphic sectional curvature c_1 and $\overline{M}^n$ be a Kaehlerian manifold of complex dimension n with constant holomorphic sectional curvature c_2. We denote by J_m and J_n almost complex structures of $\overline{M}^m$ and $\overline{M}^n$ respectively. We consider the kaeherian product $\overline{M} = \overline{M}^m \times \overline{M}^n$ with product metric g and put

$$JX = J_m\overline{P}X + J_n\overline{Q}X$$

for any vector field X on $\overline{M}$, where $\overline{P}$ and $\overline{Q}$ denote the projection operators. Then we have

$$J_m\overline{P} = \overline{P}J, \quad J_n\overline{Q} = \overline{Q}J, \quad FJ = JF, \quad F^2 = I,$$
$$J^2 = -I, \quad g(JX, JY) = g(X, Y), \quad \overline{\nabla}_X J = 0,$$

where $F = \overline{P} - \overline{Q}$ is a almost product structure on $\overline{M}$. Then the Riemannian curvature tensor $\overline{R}$ is given by

$$\begin{aligned}
\overline{R}(X,Y)Z &= \frac{c_1+c_2}{16}[g(Y,Z)X - g(X,Z)Y + g(JY,Z)JX \\
&\quad -g(JX,Z)JY + 2g(X,JY)JZ + 2g(FY,Z)FX \\
&\quad -g(FX,Z)FY + g(FJY,Z)FJX \\
&\quad -g(FJX,Z)FJY + 2g(FX,JY)FJZ] \\
&\quad +\frac{c_1-c_2}{16}[g(FY,Z)X - g(FX,Z)Y + g(Y,Z)FX \\
&\quad -g(X,Z)FY + g(FJY,Z)JX - g(FJX,Z)JY \\
&\quad +g(JY,Z)FJX - g(JX,Z)FJY \\
&\quad +2g(FX,JY)JZ + 2g(X,JY)JFZ]
\end{aligned} \tag{2.1}$$

for any vector fields X, Y and Z on $\overline{M}$.

Let M be a real l-dimensional submanifold of $\overline{M}$ with induced metric. We denote by $\overline{\nabla}$ the Levi-Civita connection in $\overline{M}$ and by ∇ the connection induced on M. Then the Gauss and Weingarten formulas are given respectively by

$$\overline{\nabla}_X Y = \nabla_X Y + B(X,Y), \quad \overline{\nabla}_X V = -A_V X + D_X V$$

for any vector field X and Y tangent to M and any vector field V normal to M, where D denote the normal connection.

For any vector field X tangent to M we put

$$JX = \mathcal{P}X + \mathcal{F}X, \qquad FX = fX + hX,$$

where $\mathcal{P}X$ and $\mathcal{F}X$ are the the tangential part and normal part of JX respectively, fX and hX are the tangential part and normal part of FX respectively.

For any vector field V normal to M we put

$$JV = \mathcal{T}V + \mathcal{N}V, \qquad FV = tV + sV,$$

where $\mathcal{T}V$ and $\mathcal{N}V$ are the tangential part and normal part of JV respectively, tV and sV are the tangential part and normal part of FV respectively. Then F, f, s, ht and $\mathcal{F}\mathcal{T}$ are symmetric with respect to g, $\mathcal{P}$, $\mathcal{N}$, $\mathcal{F}J$, $f\mathcal{P}+t\mathcal{F}$ and $\mathcal{F}t+\mathcal{N}s$ are skew-symmetric with respect to g, and $g(\mathcal{F}X,V)=-g(X,\mathcal{T}V)$, $g(hX,V)=g(X,tV)$, $g((h\mathcal{P}+s\mathcal{F})X,V)=-g(X,(\mathcal{P}t+\mathcal{T}s)V)$. We also have

$$\mathcal{P}^2=-I-\mathcal{T}\mathcal{F},\quad \mathcal{F}\mathcal{P}+\mathcal{N}\mathcal{F}=0,\quad \mathcal{P}\mathcal{T}+\mathcal{T}\mathcal{N}=0,\quad \mathcal{N}^2=-I-\mathcal{F}\mathcal{T}. \tag{2.2}$$

$$f^2=I-th,\quad hf+sh=0,\quad ft+ts=0\quad s^2=I-ht, \tag{2.3}$$

$$f\mathcal{T}+t\mathcal{N}=\mathcal{P}t+\mathcal{T}s,\quad h\mathcal{P}+s\mathcal{F}=\mathcal{F}f+\mathcal{N}h, \tag{2.4}$$

$$f\mathcal{P}+t\mathcal{F}=\mathcal{P}f+\mathcal{T}h,\quad h\mathcal{T}+s\mathcal{N}=\mathcal{F}t+\mathcal{N}s, \tag{2.5}$$

$$\begin{aligned} FJX=JFX &= (f\mathcal{P}+t\mathcal{F})X+(h\mathcal{P}+s\mathcal{F})X \\ &= (\mathcal{P}f+\mathcal{T}h)X+(\mathcal{F}f+\mathcal{N}h)X, \end{aligned} \tag{2.6}$$

$$\begin{aligned} FJV=JFV &= (\mathcal{P}t+\mathcal{T}s)V+(\mathcal{F}t+\mathcal{N}s)V \\ &= (f\mathcal{T}+t\mathcal{N})V+(h\mathcal{T}+s\mathcal{N})V. \end{aligned} \tag{2.7}$$

Additionally, we have the following relations:

$$tr(f\mathcal{P})=tr(s\mathcal{N})=tr(h\mathcal{T})=tr(\mathcal{F}t)=0,\quad tr[(h\mathcal{T})^2]=tr[(\mathcal{F}t)^2]$$

Now we define the covariant derivatives of $\mathcal{P}$, $\mathcal{F}$, $\mathcal{T}$ and $\mathcal{N}$ respectively by

$$(\nabla_X\mathcal{P})Y=\nabla_X(\mathcal{P}Y)-\mathcal{P}(\nabla_XY),\ (\nabla_X\mathcal{F})Y=D_X(\mathcal{F}Y)-\mathcal{F}(\nabla_XY),$$

$$(\nabla_X\mathcal{T})V=\nabla_X(\mathcal{T}V)-\mathcal{T}(D_XV),\ (\nabla_X\mathcal{N})V=D_X(\mathcal{N}V)-\mathcal{N}(D_XV).$$

Then we have

$$(\nabla_X\mathcal{P})Y=A_{\mathcal{F}Y}X+\mathcal{T}B(X,Y),\ (\nabla_X\mathcal{F})Y=-B(X,\mathcal{P}Y)+\mathcal{N}B(X,Y),$$

$$(\nabla_X\mathcal{T})V=-\mathcal{P}A_VX+A_{\mathcal{N}V}X,\ (\nabla_X\mathcal{N})V=-\mathcal{F}A_VX-B(X,\mathcal{T}V).$$

Because $\overline{M}$ is equipped with product metric we know that $\overline{\nabla}F=\overline{\nabla}\,\overline{P}=\overline{\nabla}\,\overline{Q}=0$ (cf [6]). Next we define the covariant derivatives of f, h, t, and s by

$$(\nabla_Xf)Y=\nabla_X(fY)-f(\nabla_XY),\quad (\nabla_Xh)Y=D_X(hY)-h(\nabla_XY),$$

$$(\nabla_Xt)V=\nabla_X(tV)-t(D_XV),\quad (\nabla_Xs)V=D_X(sV)-s(D_XV).$$

Then we have

$$(\nabla_Xf)Y=A_{hY}X+tB(x,y),\quad (\nabla_Xh)Y=sB(X,Y)-B(X,fY),$$

$$(\nabla_X t)V = A_{sV}X - fA_VX, \quad (\nabla_X s)V = -B(X,tV) - hA_VX$$

Denote by R the Riemannian curvature tensor of M, then the equation of Gauss is given by

$$R(X,Y)Z = (\overline{R}(X,Y)Z)^\top + A_{B(Y,Z)}X - A_{B(X,Z)}Y,$$

that is

$$\begin{aligned} R(X,Y)Z &= \frac{c_1+c_2}{16}[g(Y,Z)X - g(X,Z)Y + g(\mathcal{P}Y,Z)\mathcal{P}X - g(\mathcal{P}X,Z)\mathcal{P}Y \\ &\quad +2g(X,\mathcal{P}Y)\mathcal{P}Z + 2g(fY,Z)fX - g(fX,Z)fY \\ &\quad +g((f\mathcal{P}+t\mathcal{F})Y,Z)(f\mathcal{P}+t\mathcal{F})X - g((f\mathcal{P}+t\mathcal{F})X,Z)(f\mathcal{P}+t\mathcal{F})Y \\ &\quad +2g(X,(f\mathcal{P}+t\mathcal{F})Y)(f\mathcal{P}+t\mathcal{F})Z] \\ &\quad +\frac{c_1-c_2}{16}[g(fY,Z)X - g(fX,Z)Y + g(Y,Z)fX - g(X,Z)fY \\ &\quad +g((f\mathcal{P}+t\mathcal{F})Y,Z)\mathcal{P}X - g((f\mathcal{P}+t\mathcal{F})X,Z)\mathcal{P}Y \\ &\quad +g(\mathcal{P}Y,Z)(f\mathcal{P}+t\mathcal{F})X - g(\mathcal{P}X,Z)(f\mathcal{P}+t\mathcal{F})Y \\ &\quad +2g(X,(f\mathcal{P}+t\mathcal{F})Y)\mathcal{P}Z + 2g(X,\mathcal{P}Y)(f\mathcal{P}+t\mathcal{F})Z] \\ &\quad +A_{B(Y,Z)}X - A_{B(X,Z)}Y. \end{aligned} \tag{2.8}$$

The equation of Codazzi is given by

$$(\nabla_X B)(Y,Z) - (\nabla_Y B)(X,Z) = (\overline{R}(X,Y)Z)^\perp,$$

that is

$$\begin{aligned} &(\nabla_X B)(Y,Z) - (\nabla_Y B)(X,Z) \\ =\ &\frac{c_1+c_2}{16}[g(\mathcal{P}Y,Z)\mathcal{F}X - g(\mathcal{P}X,Z)\mathcal{F}Y + 2g(X,\mathcal{P}Y)\mathcal{F}Z + 2g(fY,Z)hX \\ &-g(fX,Z)hY + g((f\mathcal{P}+t\mathcal{F})Y,Z)(h\mathcal{P}+s\mathcal{F})X \\ &-g((f\mathcal{P}+t\mathcal{F})X,Z)(h\mathcal{P}+s\mathcal{F})Y + 2g(X,(f\mathcal{P}+t\mathcal{F})Y)(h\mathcal{P}+s\mathcal{F})Z] \\ &+\frac{c_1-c_2}{16}[g(Y,Z)hX - g(X,Z)hY + g((f\mathcal{P}+t\mathcal{F})Y,Z)\mathcal{F}X \\ &-g((f\mathcal{P}+t\mathcal{F})X,Z)\mathcal{F}Y + g(\mathcal{P}Y,Z)(h\mathcal{P}+s\mathcal{F})X \\ &-g(\mathcal{P}X,Z)(h\mathcal{P}+s\mathcal{F})Y + 2g(X,(f\mathcal{P}+t\mathcal{F})Y)\mathcal{F}Z \\ &+2g(X,\mathcal{P}Y)(h\mathcal{P}+s\mathcal{F})Z] \end{aligned} \tag{2.9}$$

The equation of Ricci is given by

$$R^\perp(X,Y)U = (\overline{R}(X,Y)U)^\perp + B(X,A_UY) - B(Y,A_UX)$$

or equivalently

$$g(R^\perp(X,Y)U,V) + g([A_V,A_U]X,Y) = g((\overline{R}(X,Y)U)^\perp,V),$$

where $[A_V, A_U] = A_V A_U - A_U A_V$. That is

$$\begin{aligned}
& g(R^{\perp}(X,Y)U,V) + g([A_V, A_U]X, Y) \\
= \; & \frac{c_1+c_2}{16}[g(\mathcal{F}Y,U)g(\mathcal{F}X,V) - g(\mathcal{F}X,U)g(\mathcal{F}Y,V) \\
& +2g(X,\mathcal{P}Y)g(\mathcal{N}U,V) + 2g(hY,U)g(hX,V) - g(hX,U)g(hY,V) \\
& +g((h\mathcal{P}+s\mathcal{F})Y,U)g((h\mathcal{P}+s\mathcal{F})X,V) \\
& -g((h\mathcal{P}+s\mathcal{F})X,U)g((h\mathcal{P}+s\mathcal{F})Y,V) \qquad (2.10) \\
& +2g(X,(f\mathcal{P}+t\mathcal{F})Y)((\mathcal{F}t+\mathcal{N}s)U,V)] \\
& +\frac{c_1-c_2}{16}[g((h\mathcal{P}+s\mathcal{F})Y,U)g(\mathcal{F}X,V) - g((h\mathcal{P}+s\mathcal{F})X,U)g(\mathcal{F}Y,V) \\
& +g(\mathcal{F}Y,U)g((h\mathcal{P}+s\mathcal{F})X,V) - g(\mathcal{F}X,U)g((h\mathcal{P}+s\mathcal{F})Y,V) \\
& +2g(X,(f\mathcal{P}+t\mathcal{F})Y)g(\mathcal{N}U,V) + 2g(X,\mathcal{P}Y)g((\mathcal{F}t+\mathcal{N}s)U,V)]
\end{aligned}$$

In the following, we denote by A_α the second fundamental form in the direction of v_α, where $\{v_1, \cdots, v_p\}$ is an orthonormal basis for $T_x(M)^{\perp}$, $P = 2m + 2n - l$. We denote by $|\cdot|$ the length of the tensor.

Lemma 2.1 Let M be an l-dimensional submanifold in $\overline{M}^m \times \overline{M}^n$. If the normal connection of M is flat, then

$$\begin{aligned}
& \sum_{\alpha,\beta} g([A_\beta, A_\alpha]\mathcal{T}v_\beta, \mathcal{T}v_\alpha) \\
= \; & \sum_{\alpha,\beta}[g(A_\alpha \mathcal{T}v_\beta, A_\beta \mathcal{T}v_\alpha) - g(A_\alpha \mathcal{T}v_\alpha, A_\beta \mathcal{T}v_\beta)] \\
= \; & \frac{c_1+c_2}{16}\Big\{[tr(\mathcal{F}\mathcal{T})]^2 - tr[(\mathcal{F}\mathcal{T})^2] + 2tr(\mathcal{N}\mathcal{F}\mathcal{P}\mathcal{T}) - tr[(\mathcal{F}t)^2] \qquad (2.11) \\
& +[tr(h\mathcal{P}+s\mathcal{F})\mathcal{T}]^2 - tr[(h\mathcal{P}+s\mathcal{F})\mathcal{T}]^2 + 2tr[(\mathcal{F}t+\mathcal{N}s)\mathcal{F}(f\mathcal{P}+t\mathcal{F})\mathcal{T}]\Big\} \\
& +2\frac{c_1-c_2}{16}\Big\{tr(\mathcal{F}\mathcal{T})tr((h\mathcal{P}+s\mathcal{F})\mathcal{T}) - tr(\mathcal{F}\mathcal{T}(h\mathcal{P}+s\mathcal{F})\mathcal{T}) \\
& +tr(\mathcal{N}\mathcal{F}(f\mathcal{P}+t\mathcal{F})\mathcal{T}) + tr(\mathcal{F}\mathcal{P}\mathcal{T}(\mathcal{F}t+\mathcal{N}s))\Big\}
\end{aligned}$$

where

$$-tr[(\mathcal{F}\mathcal{T})^2] = -\sum_{\alpha} g(\mathcal{F}\mathcal{T}v_\alpha, \mathcal{F}\mathcal{T}v_\alpha) \le 0 \qquad (2.12)$$

$$\begin{aligned}
& \sum_{i,\alpha} g([A_{\mathcal{N}v_\alpha}, A_\alpha]e_i, \mathcal{P}e_i) = 2\sum_{\alpha} tr(A_\alpha A_{\mathcal{N}v_\alpha}\mathcal{P}) \\
= \; & \frac{c_1+c_2}{16}\Big\{-2tr(\mathcal{T}\mathcal{N}\mathcal{F}\mathcal{P}) - 2tr(\mathcal{P}^2)tr(\mathcal{N}^2) + 3tr(\mathcal{N}h\mathcal{P}t) \qquad (2.13) \\
& -2tr[(\mathcal{P}t+\mathcal{T}s)\mathcal{N}(h\mathcal{P}+s\mathcal{F})\mathcal{P}] - 2tr[(f\mathcal{P}+t\mathcal{F})\mathcal{P}]\cdot tr[(\mathcal{F}t+\mathcal{N}s)\mathcal{N}]\Big\} \\
& -2\frac{c_1-c_2}{16}\Big\{2tr(\mathcal{P}\mathcal{T}\mathcal{N}(h\mathcal{P}+s\mathcal{F})) + tr(\mathcal{N}^2)tr((f\mathcal{P}+t\mathcal{F})\mathcal{P}) \\
& +tr(\mathcal{P}^2)tr((\mathcal{F}t+\mathcal{N}s)\mathcal{N})\Big\}
\end{aligned}$$

$$\sum_{\alpha,\beta}[g(A_\alpha tv_\beta, A_\beta tv_\alpha) - g(A_\alpha tv_\alpha, A_\beta tv_\beta)] = g([A_\alpha, A_\beta]tv_\alpha, tv_\beta) \tag{2.14}$$

$$\begin{aligned}
= \;& \frac{c_1+c_2}{16}\Big[(tr(\mathcal{F}t))^2 - tr((\mathcal{F}t)^2) - 2tr(\mathcal{N}h\mathcal{P}t) + 2(tr(ht))^2 - tr((ht)^2) \\
& +(tr((h\mathcal{P}+s\mathcal{F})t))^2 - tr(((h\mathcal{P}+s\mathcal{F})t)^2) - 2tr((\mathcal{F}t+\mathcal{N}s)h(f\mathcal{P}+t\mathcal{F})t)\Big] \\
& +2\frac{c_1-c_2}{16}\Big[-tr((h\mathcal{P}+s\mathcal{F})t\mathcal{F}t) - tr((f\mathcal{P}+t\mathcal{F})t\mathcal{N}h) - tr(h\mathcal{P}t(\mathcal{F}t+\mathcal{N}s))\Big]
\end{aligned}$$

where

$$-tr((ht)^2) = -\sum_\alpha g(htv_\alpha, htv_\alpha) \le 0 \tag{2.15}$$

$$\begin{aligned}
& g([A_{(\mathcal{F}t+\mathcal{N}s)v_\alpha}, A_\alpha]e_i, (f\mathcal{P}+t\mathcal{F})e_i) \\
= \;& 2tr(A_\alpha A_{(\mathcal{F}t+\mathcal{N}s)v_\alpha}(f\mathcal{P}+t\mathcal{F})) \qquad (2.16) \\
= \;& \frac{c_1+c_2}{16}\Big\{-2tr((f\mathcal{P}+t\mathcal{F})\mathcal{T}(\mathcal{F}t+\mathcal{N}s)\mathcal{F}) - 2tr(\mathcal{P}(f\mathcal{P}+t\mathcal{F}))tr(\mathcal{N}(\mathcal{F}t+\mathcal{N}s)) \\
& +3tr(t(\mathcal{F}t+\mathcal{N}s)h(f\mathcal{P}+t\mathcal{F})) - 2tr((\mathcal{P}t+\mathcal{T}s)(\mathcal{F}t+\mathcal{N}s)(h\mathcal{P}+s\mathcal{F})(f\mathcal{P}+t\mathcal{F})) \\
& -2tr((f\mathcal{P}+t\mathcal{F})^2)\cdot tr((\mathcal{F}t+\mathcal{N}s)^2)\Big\} \\
& -2\frac{c_1-c_2}{16}\Big\{tr(\mathcal{T}(\mathcal{F}t+\mathcal{N}s)(h\mathcal{P}+s\mathcal{F})(f\mathcal{P}+t\mathcal{F})) \\
& +tr((\mathcal{P}t+\mathcal{T}s)(\mathcal{F}t+\mathcal{N}s)\mathcal{F}(f\mathcal{P}+t\mathcal{F})) \\
& +tr((f\mathcal{P}+t\mathcal{F})^2)tr(\mathcal{N}(\mathcal{F}t+\mathcal{N}s)) + tr((\mathcal{F}t+\mathcal{N}s)^2)tr(\mathcal{P}(f\mathcal{P}+t\mathcal{F}))\Big\}
\end{aligned}$$

where

$$\begin{aligned}
& -2tr((f\mathcal{P}+t\mathcal{F})^2)\cdot tr((\mathcal{F}t+\mathcal{N}s)^2) \\
= \;& -2\sum_{i,\alpha} g((f\mathcal{P}+t\mathcal{F})e_i, (f\mathcal{P}+t\mathcal{F})e_i)\cdot g((\mathcal{F}t+\mathcal{N}s)v_\alpha, (\mathcal{F}t+\mathcal{N}s)v_\alpha) \\
\le \;& 0 \qquad (2.17)
\end{aligned}$$

$$\begin{aligned}
& \sum_{\alpha,\beta}\Big[g(A_\alpha(\mathcal{P}t+\mathcal{T}s)v_\beta, A_\beta(\mathcal{P}t+\mathcal{T}s)v_\alpha) - g(A_\alpha(\mathcal{P}t+\mathcal{T}s)v_\alpha, A_\beta(\mathcal{P}t+\mathcal{T}s)v_\beta)\Big] \\
= \;& \sum_{\alpha,\beta} g([A_\alpha, A_\beta](\mathcal{P}t+\mathcal{T}s)v_\alpha, (\mathcal{P}t+\mathcal{T}s)v_\beta) \qquad (2.18) \\
= \;& \frac{c_1+c_2}{16}\Big\{[tr(\mathcal{F}(\mathcal{P}t+\mathcal{T}s))]^2 - tr[(\mathcal{F}(\mathcal{P}t+\mathcal{T}s))^2] \\
& +2tr(\mathcal{N}(h\mathcal{P}+s\mathcal{F})\mathcal{P}(\mathcal{P}t+\mathcal{T}s)) + 2[tr(h(\mathcal{P}t+\mathcal{T}s))]^2 - tr[(h(\mathcal{P}t+\mathcal{T}s))^2] \\
& +[tr((h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s))]^2 - tr[((h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s))^2] \\
& +2tr((\mathcal{F}t+\mathcal{N}s)(h\mathcal{P}+s\mathcal{F})(f\mathcal{P}+t\mathcal{F})(\mathcal{P}t+\mathcal{T}s))\Big\} \\
& +2\frac{c_1-c_2}{16}\Big\{tr(\mathcal{F}(\mathcal{P}t+\mathcal{T}s))\cdot tr((h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s)) \\
& -tr(\mathcal{F}(\mathcal{P}t+\mathcal{T}s)(h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s)) + tr(\mathcal{N}(h\mathcal{P}+s\mathcal{F})(f\mathcal{P}+t\mathcal{F})(\mathcal{P}t+\mathcal{T}s)) \\
& +tr((\mathcal{F}t+\mathcal{N}s)(h\mathcal{P}+s\mathcal{F})\mathcal{P}(\mathcal{P}t+\mathcal{T}s))\Big\}
\end{aligned}$$

where

$$\begin{aligned} & -tr[((h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s))^2] \\ = & -\sum_{\alpha} g((h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s)v_\alpha,(h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s)v_\alpha) \\ \leq & \ 0 \end{aligned} \tag{2.19}$$

$$\begin{aligned} & g([A_{(\mathcal{F}t+\mathcal{N}s)v_\alpha},A_\alpha]e_i,\mathcal{P}e_i) = 2tr(A_\alpha A_{(\mathcal{F}t+\mathcal{N}s)v_\alpha}\mathcal{P}) \\ = & \frac{c_1+c_2}{16}\Big\{-2tr(\mathcal{F}\mathcal{P}\mathcal{T}(\mathcal{F}t+\mathcal{N}s))-2tr(\mathcal{P}^2)tr(\mathcal{N}(\mathcal{F}t+\mathcal{N}s)) \\ & +3tr(h\mathcal{P}t(\mathcal{F}t+\mathcal{N}s))-2tr((\mathcal{P}t+\mathcal{T}s)(\mathcal{F}t+\mathcal{N}s)(h\mathcal{P}+s\mathcal{F})\mathcal{P}) \\ & -2tr((f\mathcal{P}+t\mathcal{F})\mathcal{P})\cdot tr((\mathcal{F}t+\mathcal{N}s)^2)\Big\} \\ & +\frac{c_1-c_2}{16}\Big\{-4tr(\mathcal{P}\mathcal{T}(\mathcal{F}t+\mathcal{N}s)(h\mathcal{P}+s\mathcal{F})) \\ & -2tr((f\mathcal{P}+t\mathcal{F})\mathcal{P})\cdot tr(\mathcal{N}(\mathcal{F}t+\mathcal{N}s))-2tr(\mathcal{P}^2)\cdot tr((\mathcal{F}t+\mathcal{N}s)^2)\Big\} \end{aligned} \tag{2.20}$$

where

$$\begin{aligned} & -2tr(\mathcal{P}^2)\cdot tr((\mathcal{F}t+\mathcal{N}s)^2) \\ = & -2g(\mathcal{P}e_i,\mathcal{P}e_i)\cdot g((\mathcal{F}t+\mathcal{N}s)v_\alpha,(\mathcal{F}t+\mathcal{N}s)v_\alpha)\leq 0 \end{aligned} \tag{2.21}$$

$$\begin{aligned} & \sum_{\alpha,\beta}\Big[g(A_\alpha\mathcal{T}v_\beta,A_\beta(\mathcal{P}t+\mathcal{T}s)v_\alpha)-g(A_\alpha\mathcal{T}v_\alpha,A_\beta(\mathcal{P}t+\mathcal{T}s)v_\beta)\Big] \\ = & \sum_{\alpha,\beta} g([A_\alpha,A_\beta](\mathcal{P}t+\mathcal{T}s)v_\alpha,\mathcal{T}v_\beta) \\ = & \frac{c_1+c_2}{16}\Big\{tr(\mathcal{F}\mathcal{T})\cdot tr(\mathcal{F}(\mathcal{P}t+\mathcal{T}s))-tr(\mathcal{F}\mathcal{T}\mathcal{F}(\mathcal{P}t+\mathcal{T}s)) \\ & +2tr(\mathcal{N}\mathcal{F}\mathcal{P}(\mathcal{P}t+\mathcal{T}s))+2tr(h\mathcal{T})\cdot tr(h(\mathcal{P}t+\mathcal{T}s)) \\ & -tr(h\mathcal{T}h(\mathcal{P}t+\mathcal{T}s))+tr((h\mathcal{P}+s\mathcal{F})\mathcal{T})\cdot tr((h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s)) \\ & -tr((h\mathcal{P}+s\mathcal{F})\mathcal{T}(h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s)) \\ & +2tr((\mathcal{F}t+\mathcal{N}s)((h\mathcal{P}+s\mathcal{F})(f\mathcal{P}+t\mathcal{F})\mathcal{T})\Big\} \\ & +\frac{c_1-c_2}{16}\Big\{(tr(\mathcal{F}(\mathcal{P}t+\mathcal{T}s)))^2-tr(\mathcal{T}\mathcal{F}(\mathcal{P}t+\mathcal{T}s)(h\mathcal{P}+s\mathcal{F})) \\ & -tr(\mathcal{F}\mathcal{T}(h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s))+tr(\mathcal{F}\mathcal{T})\cdot tr((h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s)) \\ & +2tr(\mathcal{T}\mathcal{N}(h\mathcal{P}+s\mathcal{F})(f\mathcal{P}+t\mathcal{F}))+2tr((\mathcal{F}t+\mathcal{N}s)(h\mathcal{P}+s\mathcal{F})\mathcal{P}\mathcal{T})\Big\} \end{aligned} \tag{2.22}$$

where

$$\begin{aligned} & -tr(\mathcal{T}\mathcal{F}(\mathcal{P}t+\mathcal{T}s)(h\mathcal{P}+s\mathcal{F})) \\ = & -\sum_{\alpha} g(\mathcal{F}(\mathcal{P}t+\mathcal{T}s)v_\alpha,\mathcal{F}(\mathcal{P}t+\mathcal{T}s)v_\alpha) \\ \leq & \ 0 \end{aligned} \tag{2.23}$$

and

$$\begin{aligned} & tr(\mathcal{F}\mathcal{T})\cdot tr((h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s)) \\ = & \sum_{\alpha,\beta} g(\mathcal{T}v_\alpha,\mathcal{T}v_\alpha)\cdot g((\mathcal{P}t+\mathcal{T}s)v_\beta,(\mathcal{P}t+\mathcal{T}s)v_\beta) \\ \geq & 0 \end{aligned} \tag{2.24}$$

$$\begin{aligned} & g([A_{\mathcal{N}v_\alpha},A_\alpha]e_i,(f\mathcal{P}+t\mathcal{F})e_i) = 2tr(A_\alpha A_{\mathcal{N}v_\alpha}(f\mathcal{P}+t\mathcal{F})) \qquad (2.25) \\ = & \frac{c_1+c_2}{16}\Big\{-2tr(\mathcal{T}\mathcal{N}\mathcal{F}(f\mathcal{P}+t\mathcal{F}))-2tr(\mathcal{P}(f\mathcal{P}+t\mathcal{F}))\cdot tr(\mathcal{N}^2) \\ & +3tr(t\mathcal{N}h(f\mathcal{P}+t\mathcal{F}))-2tr((\mathcal{P}t+\mathcal{T}s)\mathcal{N}(h\mathcal{P}+s\mathcal{F})(f\mathcal{P}+t\mathcal{F})) \\ & -2tr((f\mathcal{P}+t\mathcal{F})^2)\cdot tr(\mathcal{N}(\mathcal{F}t+\mathcal{N}s))\Big\} \\ & +\frac{c_1-c_2}{16}\Big\{-2tr(\mathcal{T}\mathcal{N}(h\mathcal{P}+s\mathcal{F})(f\mathcal{P}+t\mathcal{F}))-2tr(\mathcal{N}\mathcal{F}(f\mathcal{P}+t\mathcal{F})(\mathcal{P}t+\mathcal{T}s)) \\ & -2tr((f\mathcal{P}+t\mathcal{F})^2)\cdot tr(\mathcal{N}^2)-2tr(\mathcal{P}(f\mathcal{P}+t\mathcal{F}))\cdot tr(\mathcal{N}(\mathcal{F}t+\mathcal{N}s))\Big\} \end{aligned}$$

where

$$\begin{aligned} & -2tr((f\mathcal{P}+t\mathcal{F})^2)\cdot tr(\mathcal{N}^2) \\ = & -2g((f\mathcal{P}+t\mathcal{F})e_i,(f\mathcal{P}+t\mathcal{F})e_i)\cdot g(\mathcal{N}v_\alpha,\mathcal{N}v_\alpha) \\ \leq & 0 \end{aligned} \tag{2.26}$$

3 Minimal submanifolds with flat normal connection

In this section, we give the Simons' type integral formula for a compact minimal submanifold M in $\overline{M}^m\times\overline{M}^n$. According to [4], we have to compute the following terms:

$$\begin{aligned} & \sum_i(\overline{\nabla}_X(\overline{R}(e_i,Y)e_i)^\perp)^\perp \\ = & \sum_i(\overline{R}(B(X,e_i),Y)e_i)^\perp+\sum_i(\overline{R}(e_i,B(X,Y))e_i)^\perp \\ & +\sum_i(\overline{R}(e_i,Y)B(X,e_i))^\perp-\sum_i B(X,(\overline{R}(e_i,Y)e_i)^\top) \end{aligned} \tag{3.27}$$

where

$$
\begin{aligned}
&\sum_i (\overline{R}(B(X,e_i),Y)e_i)^{\perp} \\
= \;& \frac{c_1+c_2}{16}\Big[B(X,Y)+\mathcal{N}B(X,\mathcal{P}Y) \\
&+\sum_i g(B(X,e_i),\mathcal{F}e_i)\mathcal{F}Y+2\sum_i g(B(X,e_i),\mathcal{F}Y)\mathcal{F}e_i \\
&+2sB(X,fY)-\sum_i g(B(X,e_i),he_i)hY+(\mathcal{F}t+\mathcal{N}s)B(X,(f\mathcal{P}+t\mathcal{F})Y) \\
&+\sum_i g(B(X,e_i),(h\mathcal{P}+s\mathcal{F})e_i)(h\mathcal{P}+s\mathcal{F})Y \qquad (3.28)\\
&+2\sum_i g(B(X,e_i),(h\mathcal{P}+s\mathcal{F})Y)(h\mathcal{P}+s\mathcal{F})e_i\Big] \\
&+\frac{c_1-c_2}{16}\Big[B(X,fY)+sB(X,Y)+\mathcal{N}B(X,(f\mathcal{P}+t\mathcal{F})Y) \\
&+g(B(X,e_i),(h\mathcal{P}+s\mathcal{F})e_i)\mathcal{F}Y+(\mathcal{F}t+\mathcal{N}s)B(X,\mathcal{P}Y) \\
&+g(B(X,e_i),\mathcal{F}e_i)(h\mathcal{P}+s\mathcal{F})Y+2g(B(X,e_i),(h\mathcal{P}+s\mathcal{F})Y)\mathcal{F}e_i \\
&+2g(B(X,e_i),\mathcal{F}Y)(h\mathcal{P}+s\mathcal{F})e_i\Big]
\end{aligned}
$$

$$
\begin{aligned}
&\sum_i (\overline{R}(e_i,B(X,Y))e_i)^{\perp} \\
= \;& \frac{c_1+c_2}{16}\Big[-l\cdot B(X,Y)+3\mathcal{F}\mathcal{T}B(X,Y)+2htB(X,Y) \qquad (3.29)\\
&-tr(f)sB(X,Y)+3(h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s)B(X,Y)\Big] \\
&+\frac{c_1-c_2}{16}\Big[-tr(f)B(X,Y)-l\cdot sB(X,Y) \\
&+3\mathcal{F}(\mathcal{P}t+\mathcal{T}s)B(X,Y)+3(h\mathcal{P}+s\mathcal{F})\mathcal{T}B(X,Y)\Big]
\end{aligned}
$$

$$
\begin{aligned}
&\sum_i (\overline{R}(e_i,Y)B(X,e_i))^{\perp} \\
= \;& \frac{c_1+c_2}{16}\Big[\sum_i g(\mathcal{F}Y,B(X,e_i))\mathcal{F}e_i - \sum_i g(\mathcal{F}e_i,B(X,e_i))\mathcal{F}Y \\
&+2\mathcal{N}B(X,\mathcal{P}Y)+2\sum_i g(hY,B(X,e_i))he_i - \sum_i g(he_i,B(X,e_i))hY \\
&+\sum_i g((h\mathcal{P}+s\mathcal{F})Y,B(X,e_i))(h\mathcal{P}+s\mathcal{F})e_i \\
&-\sum_i g((h\mathcal{P}+s\mathcal{F})e_i,B(X,e_i))(h\mathcal{P}+s\mathcal{F})Y \\
&+2(\mathcal{F}t+\mathcal{N}s)B(X,(f\mathcal{P}+t\mathcal{F})Y)\Big] \\
&+\frac{c_1-c_2}{16}\Big[g((h\mathcal{P}+s\mathcal{F})Y,B(X,e_i))\mathcal{F}e_i - g((h\mathcal{P}+s\mathcal{F})e_i,B(X,e_i))\mathcal{F}Y \\
&+g(\mathcal{F}Y,B(X,e_i))(h\mathcal{P}+s\mathcal{F})e_i - g(\mathcal{F}e_i,B(X,e_i))(h\mathcal{P}+s\mathcal{F})Y \\
&+2\mathcal{N}B(X,(f\mathcal{P}+t\mathcal{F})Y)+2(\mathcal{F}t+\mathcal{N}s)B(X,\mathcal{P}Y)\Big]
\end{aligned} \tag{3.30}
$$

$$
\begin{aligned}
&\sum_i (\overline{R}(e_i,Y)e_i)^{\top} \\
= \;& \frac{c_1+c_2}{16}\Big[(1-l)Y+3\mathcal{P}^2Y+2f^2Y-tr(f)\cdot fY+3(f\mathcal{P}+t\mathcal{F})^2Y\Big] \\
&+\frac{c_1-c_2}{16}\Big[(2-l)fY-tr(f)Y+3\mathcal{P}(f\mathcal{P}+t\mathcal{F})Y+3(f\mathcal{P}+t\mathcal{F})\mathcal{P}Y\Big]
\end{aligned} \tag{3.31}
$$

Therefore

$$
\begin{aligned}
&-\sum_i B(X,(\overline{R}(e_i,Y)e_i)^{\top}) \\
= \;& -\frac{c_1+c_2}{16}\Big[(1-l)B(X,Y)+3B(X,\mathcal{P}^2Y)+2B(X,f^2Y) \\
&-tr(f)B(X,fY)+3B(X,(f\mathcal{P}+t\mathcal{F})^2Y)\Big] \\
&-\frac{c_1-c_2}{16}\Big[(2-l)B(X,fY)-tr(f)B(X,Y)+3B(X,\mathcal{P}(f\mathcal{P}+t\mathcal{F})Y) \\
&+3B(X,(f\mathcal{P}+t\mathcal{F})\mathcal{P}Y)\Big]
\end{aligned} \tag{3.32}
$$

We also have to compute

$$
\begin{aligned}
&\sum_i (\overline{\nabla}_{e_i}(\overline{R}(e_i,X)Y)^{\perp})^{\perp} \\
= \;& \sum_i (\overline{R}(e_i,B(e_i,X))Y)^{\perp}+\sum_i (\overline{R}(e_i,X)B(e_i,Y))^{\perp}-\sum_i B(e_i,(\overline{R}(e_i,X)Y)^{\top})
\end{aligned} \tag{3.33}
$$

where

$$
\begin{aligned}
&\sum_i (\overline{R}(e_i, B(e_i,X))Y)^\perp \\
= \;& \frac{c_1+c_2}{16}\Big[- B(X,Y) - sB(X,fY) + \mathcal{N}B(X,\mathcal{P}Y) \\
&- \sum_i g(B(e_i,X),\mathcal{F}Y)\mathcal{F}e_i - 2\sum_i g(B(e_i,X),\mathcal{F}e_i)\mathcal{F}Y \\
&+2\sum_i g(B(e_i,X),hY)he_i + (\mathcal{F}t+\mathcal{N}s)B((f\mathcal{P}+t\mathcal{F})Y,X) \\
&- \sum_i g(B(e_i,X),(h\mathcal{P}+s\mathcal{F})Y)(h\mathcal{P}+s\mathcal{F})e_i \\
&-2\sum_i g(B(e_i,X),(h\mathcal{P}+s\mathcal{F})e_i)(h\mathcal{P}+s\mathcal{F})Y\Big] \\
&+\frac{c_1-c_2}{16}\Big[- B(X,fY) - hB(X,Y) + \mathcal{N}B(X,(f\mathcal{P}+t\mathcal{F})Y) \\
&+(\mathcal{F}t+\mathcal{N}s)B(X,\mathcal{P}Y) \\
&-g(B(e_i,X),(h\mathcal{P}+s\mathcal{F})Y)\mathcal{F}e_i - 2g(B(e_i,X),(h\mathcal{P}+s\mathcal{F})e_i)\mathcal{F}Y \\
&-g(B(e_i,X),\mathcal{F}Y)(h\mathcal{P}+s\mathcal{F})e_i - 2g(B(e_i,X),\mathcal{F}e_i)(h\mathcal{P}+s\mathcal{F})Y\Big]
\end{aligned}
\tag{3.34}
$$

$$
\begin{aligned}
&\sum_i (\overline{R}(e_i,X)B(Y,e_i))^\perp \\
= \;& \frac{c_1+c_2}{16}\Big[\sum_i g(\mathcal{F}X,B(Y,e_i))\mathcal{F}e_i - \sum_i g(\mathcal{F}e_i,B(Y,e_i))\mathcal{F}X \\
&+2\mathcal{N}B(Y,\mathcal{P}X) + 2\sum_i g(hX,B(Y,e_i))he_i - \sum_i g(he_i,B(Y,e_i))hX \\
&+\sum_i g((h\mathcal{P}+s\mathcal{F})X,B(Y,e_i))(h\mathcal{P}+s\mathcal{F})e_i \\
&-\sum_i g((h\mathcal{P}+s\mathcal{F})e_i,B(Y,e_i))(h\mathcal{P}+s\mathcal{F})X \\
&+2(\mathcal{F}t+\mathcal{N}s)B(Y,(f\mathcal{P}+t\mathcal{F})X)\Big] \\
&+\frac{c_1-c_2}{16}\Big[g((h\mathcal{P}+s\mathcal{F})X,B(Y,e_i))\mathcal{F}e_i - g((h\mathcal{P}+s\mathcal{F})e_i,B(Y,e_i))\mathcal{F}X \\
&+g(\mathcal{F}X,B(Y,e_i))(h\mathcal{P}+s\mathcal{F})e_i - g(\mathcal{F}e_i,B(Y,e_i))(h\mathcal{P}+s\mathcal{F})X \\
&+2\mathcal{N}B(Y,(f\mathcal{P}+t\mathcal{F})X) + 2(\mathcal{F}t+\mathcal{N}s)B(Y,\mathcal{P}X)\Big]
\end{aligned}
\tag{3.35}
$$

$$
\begin{aligned}
&\sum_i (\overline{R}(e_i,X)Y)^\top \\
= \;& \frac{c_1+c_2}{16}\sum_i \Big[g(X,Y)e_i - g(e_i,Y)X + g(\mathcal{P}X,Y)\mathcal{P}e_i + g(\mathcal{P}Y,e_i)\mathcal{P}X \\
&+2g(e_i,\mathcal{P}X)\mathcal{P}Y + 2g(fX,Y)fe_i - g(e_i,fY)fX \\
&+g((f\mathcal{P}+t\mathcal{F})X,Y)(f\mathcal{P}+t\mathcal{F})e_i \\
&+g((f\mathcal{P}+t\mathcal{F})Y,e_i)(f\mathcal{P}+t\mathcal{F})X + 2g((f\mathcal{P}+t\mathcal{F})X,e_i)(f\mathcal{P}+t\mathcal{F})Y\Big] \\
&+\frac{c_1-c_2}{16}\sum_i \Big[g(fX,Y)e_i - g(fY,e_i)X + g(X,Y)fe_i - g(Y,e_i)fX \\
&+g((f\mathcal{P}+t\mathcal{F})X,Y)\mathcal{P}e_i + g((f\mathcal{P}+t\mathcal{F})Y,e_i)\mathcal{P}X \\
&+g(\mathcal{P}X,Y)(f\mathcal{P}+t\mathcal{F})e_i + g(\mathcal{P}Y,e_i)(f\mathcal{P}+t\mathcal{F})X \\
&+2g((f\mathcal{P}+t\mathcal{F})X,e_i)\mathcal{P}Y + 2g(\mathcal{P}X,e_i)(f\mathcal{P}+t\mathcal{F})Y\Big]
\end{aligned} \tag{3.36}
$$

Therefore

$$
\begin{aligned}
&-\sum_i B(e_i,(\overline{R}(e_i,X)Y)^\top) \\
&= -\frac{c_1+c_2}{16}\Big[-B(X,Y) + 3B(\mathcal{P}X,\mathcal{P}Y) + 2g(fX,Y)B(fe_i,e_i) \\
&-B(fX,fY) + 3B((f\mathcal{P}+t\mathcal{F})X,(f\mathcal{P}+t\mathcal{F})Y)\Big] \\
&-\frac{c_1-c_2}{16}\Big[-B(X,fY) + g(X,Y)B(e_i,fe_i) - B(fX,Y) \\
&+3B(\mathcal{P}X,(f\mathcal{P}+t\mathcal{F})Y) + 3B(\mathcal{P}Y,(f\mathcal{P}+t\mathcal{F})X)\Big]
\end{aligned} \tag{3.37}
$$

Then we have

Lemma 3.1 Let M be a minimal submanifold in $\overline{M}^m \times \overline{M}^n$ with flat normal

connection, then

$$
\begin{aligned}
& g(\nabla^2 B, B) = \sum_{i,j,k} g(\nabla_{e_i}\nabla_{e_i} B(e_j,e_k), B(e_j,e_k)) \\
= & \sum_{i,j,k} \Big[g((R(e_i,e_j)B)(e_i,e_k), B(e_j,e_k)) \\
& + g((\overline{\nabla}_{e_j}(\overline{R}(e_i,e_k)e_i)^{\perp})^{\perp}, B(e_j,e_k)) + g((\overline{\nabla}_{e_i}(\overline{R}(e_i,e_j)e_k)^{\perp})^{\perp}, B(e_j,e_k)) \\
= & \sum_{i,j,k} g((R(e_i,e_j)B)(e_i,e_k), B(e_j,e_k)) \qquad (3.38) \\
& + \frac{c_1+c_2}{16}\Big\{ 3\sum_{\alpha}\Big[tr(A_\alpha A_{\mathcal{F}\mathcal{T}v_\alpha}) - 2tr(A_\alpha A_{\mathcal{N}v_\alpha}\mathcal{P}) + tr((A_\alpha\mathcal{P})^2) - tr(A_\alpha^2\mathcal{P}^2)\Big] \\
& + 3\sum_{\alpha,\beta}\Big[g(A_\alpha(\mathcal{P}t+\mathcal{T}s)v_\beta, A_\beta(\mathcal{P}t+\mathcal{T}s)v_\alpha) - g(A_\alpha(\mathcal{P}t+\mathcal{T}s)v_\alpha, A_\beta(\mathcal{P}t+\mathcal{T}s)v_\beta)\Big] \\
& + 3\sum_{\alpha,\beta}\Big[g(A_\alpha\mathcal{T}v_\beta, A_\beta\mathcal{T}v_\alpha) - g(A_\alpha\mathcal{T}v_\alpha, A_\beta\mathcal{T}v_\beta)\Big] \\
& + 3\sum_{\alpha}\Big[tr((A_\alpha(f\mathcal{P}+t\mathcal{F}))^2) - tr(A_\alpha^2(f\mathcal{P}+t\mathcal{F})^2)\Big] \\
& + 3\sum_{\alpha,\beta}\Big[g(A_\alpha tv_\beta, A_\beta tv_\alpha) - g(A_\alpha tv_\alpha, A_\beta tv_\beta)\Big] + 3\sum_{\alpha,\beta} g(A_\alpha tv_\beta, A_\beta tv_\alpha) \\
& + \sum_{\alpha}\Big[2tr(A_\alpha A_{htv_\alpha}) + tr(A_\alpha A_{sv_\alpha} f) - tr(f)tr(A_\alpha A_{sv_\alpha}) + tr(f)tr(A_\alpha^2 f) \\
& - 2tr(A_\alpha^2 f^2) - 2(tr(A_\alpha f))^2 + tr((A_\alpha f)^2) \\
& - 6tr(A_\alpha A_{(\mathcal{F}t+\mathcal{N}s)v_\alpha}(f\mathcal{P}+t\mathcal{F})) + 3tr(A_\alpha A_{(h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s)v_\alpha})\Big]\Big\}
\end{aligned}
$$

$$
\begin{aligned}
& + \frac{c_1-c_2}{16}\Big\{ l\cdot tr(A_\alpha^2 f) - l\cdot tr(A_\alpha A_{sv_\alpha}) \\
& + 6g(A_\alpha\mathcal{T}v_\beta, A_\beta(\mathcal{P}t+\mathcal{T}s)v_\alpha) - 6g(A_\alpha\mathcal{T}v_\alpha, A_\beta(\mathcal{P}t+\mathcal{T}s)v_\beta) \\
& - 6tr(A_\alpha A_{\mathcal{N}v_\alpha}(f\mathcal{P}+t\mathcal{F})) - 6tr(A_\alpha A_{(\mathcal{F}t+\mathcal{N}s)}v_\alpha\mathcal{P}) \\
& + 6tr(A_\alpha A_{(h\mathcal{P}+s\mathcal{F})\mathcal{T}v_\alpha}) - 6tr(A_\alpha^2(f\mathcal{P}+t\mathcal{F})\mathcal{P}) + 6tr(A_\alpha\mathcal{P}A_\alpha(f\mathcal{P}+t\mathcal{F}))\Big\}
\end{aligned}
$$

□

We also have the following relations:

$$\sum_{\alpha} tr(A_\alpha A_{\mathcal{F}\mathcal{T}v_\alpha}) = \sum_{\alpha}[tr(A_{\mathcal{N}v_\alpha}^2) - tr(A_\alpha^2)] \qquad (3.39)$$

$$
\begin{aligned}
\sum_{\alpha} tr(A_\alpha A_{htv_\alpha}) &= \sum_{\alpha}[tr(A_\alpha^2) - tr(A_{s^2v_\alpha})] \\
&= \sum_{\alpha}[tr(A_\alpha^2) - tr(A_{sv_\alpha}^2)] \qquad (3.40)
\end{aligned}
$$

$$\sum_{\alpha} tr(A_{s^2v_\alpha}) = \sum_{\alpha} tr(A_{sv_\alpha}^2) \qquad (3.41)$$

$$\sum_\alpha |[\mathcal{P}, A_\alpha]|^2 = 2\sum_\alpha [tr((A_\alpha \mathcal{P})^2) - tr(A_\alpha^2 \mathcal{P}^2)] \tag{3.42}$$

$$\sum_\alpha |[f, A_\alpha]|^2 = -2\sum_\alpha [tr((A_\alpha f)^2) - tr(A_\alpha^2 f^2)] \tag{3.43}$$

$$\begin{aligned} & \sum_\alpha |[f\mathcal{P} + t\mathcal{F}, A_\alpha]|^2 \\ = \; & 2\sum_\alpha [tr((A_\alpha (f\mathcal{P} + t\mathcal{F}))^2) - tr(A_\alpha^2 (f\mathcal{P} + t\mathcal{F})^2)] \end{aligned} \tag{3.44}$$

$$\sum_\alpha tr(A_\alpha A_{\mathcal{N}\alpha}) = \sum_\alpha tr(A_\alpha^2 \mathcal{P}) = \sum_\alpha tr(A_\alpha \mathcal{P}) = 0 \tag{3.45}$$

$$\sum_\alpha tr(A_\alpha (f\mathcal{P} + t\mathcal{F})) = \sum_\alpha tr(A_\alpha^2 (f\mathcal{P} + t\mathcal{F})) = 0 \tag{3.46}$$

$$\sum_\alpha tr(A_\alpha A_{(\mathcal{F}t + \mathcal{N}s)v_\alpha}) = 0 \tag{3.47}$$

$$\begin{aligned} & \sum_\alpha tr(A_\alpha A_{(h\mathcal{P} + s\mathcal{F})(\mathcal{P}t + \mathcal{T}s)v_\alpha}) \\ = \; & -\sum_{\alpha,\beta} tr(A_\alpha A_\beta) \cdot g((\mathcal{P}t + \mathcal{T}s)v_\alpha, (\mathcal{P}t + \mathcal{T}s)v_\beta) \\ = \; & -\sum_{i,j,k} \Big[g((h\mathcal{P} + s\mathcal{F})e_j, B(e_i, e_k)) \Big]^2 \leq 0 \end{aligned} \tag{3.48}$$

$$tr(A_\alpha^2 \mathcal{P}(f\mathcal{P} + t\mathcal{F})) = tr(A_\alpha^2 (f\mathcal{P} + t\mathcal{F})\mathcal{P}) \tag{3.49}$$

Now we can rewrite (3.12) as follows:

$$
\begin{aligned}
& g(\nabla^2 A, A) = g(\nabla^2 B, B) = \sum_{i,j,k} g(\nabla_{e_i}\nabla_{e_i} B(e_j, e_k), B(e_j, e_k)) \\
= \; & \sum_{i,j,\alpha} g((R(e_i, e_j)A_\alpha)e_i, A_\alpha e_j) \qquad (3.50) \\
& + \frac{c_1 + c_2}{16}\Big\{ 3\sum_\alpha \Big[tr(A^2_{\mathcal{N}v_\alpha}) - tr(A^2_\alpha) - 2tr(A_\alpha A_{\mathcal{N}v_\alpha}\mathcal{P}) + \frac{1}{2}\sum_\alpha |[\mathcal{P}, A_\alpha]|^2 \Big] \\
& + 3\frac{c_1 + c_2}{16}\Big[(tr(\mathcal{F}(\mathcal{P}t + \mathcal{T}s)))^2 - tr((\mathcal{F}(\mathcal{P}t + \mathcal{T}s))^2) \\
& + 2tr(\mathcal{N}(h\mathcal{P} + s\mathcal{F})\mathcal{P}(\mathcal{P}t + \mathcal{T}s)) + 2(tr(h(\mathcal{P}t + \mathcal{T}s)))^2 - tr((h(\mathcal{P}t + \mathcal{T}s))^2) \\
& + (tr((h\mathcal{P} + s\mathcal{F})(\mathcal{P}t + \mathcal{T}s)))^2 - tr(((h\mathcal{P} + s\mathcal{F})(\mathcal{P}t + \mathcal{T}s))^2) \\
& + 2tr((\mathcal{F}t + \mathcal{N}s)(h\mathcal{P} + s\mathcal{F})(f\mathcal{P} + t\mathcal{F})(\mathcal{P}t + \mathcal{T}s))\Big] \\
& + 6\frac{c_1 - c_2}{16}\Big[tr(\mathcal{F}(\mathcal{P}t + \mathcal{T}s)) \cdot tr((h\mathcal{P} + s\mathcal{F})(\mathcal{P}t + \mathcal{T}s)) \\
& - tr(\mathcal{F}(\mathcal{P}t + \mathcal{T}s)(h\mathcal{P} + s\mathcal{F})(\mathcal{P}t + \mathcal{T}s)) + tr(\mathcal{N}(h\mathcal{P} + s\mathcal{F})(f\mathcal{P} + t\mathcal{F})(\mathcal{P}t + \mathcal{T}s)) \\
& + tr((\mathcal{F}t + \mathcal{N}s)(h\mathcal{P} + s\mathcal{F})\mathcal{P}(\mathcal{P}t + \mathcal{T}s))\Big] \\
& + 3\frac{c_1 + c_2}{16}\Big[(tr(\mathcal{F}\mathcal{T}))^2 - tr((\mathcal{F}\mathcal{T})^2) + 2tr(\mathcal{N}\mathcal{F}\mathcal{P}\mathcal{T}) - tr((\mathcal{F}t)^2) \\
& + (tr(h\mathcal{P} + s\mathcal{F})\mathcal{T})^2 - tr(((h\mathcal{P} + s\mathcal{F})\mathcal{T})^2) + 2tr((\mathcal{F}t + \mathcal{N}s)\mathcal{F}(f\mathcal{P} + t\mathcal{F})\mathcal{T})\Big] \\
& + 6\frac{c_1 - c_2}{16}\Big[tr(\mathcal{F}\mathcal{T})tr((h\mathcal{P} + s\mathcal{F})\mathcal{T}) - tr(\mathcal{F}\mathcal{T}(h\mathcal{P} + s\mathcal{F})\mathcal{T}) \\
& + tr(\mathcal{N}\mathcal{F}(f\mathcal{P} + t\mathcal{F})\mathcal{T}) + tr(\mathcal{F}\mathcal{P}\mathcal{T}(\mathcal{F}t + \mathcal{N}s))\Big] \\
& + 3\frac{c_1 + c_2}{16}\Big[- tr((\mathcal{F}t)^2) - 2tr(\mathcal{N}h\mathcal{P}t) + 2(tr(ht))^2 - tr((ht)^2)
\end{aligned}
$$

$$
\begin{aligned}
&+(tr((h\mathcal{P}+s\mathcal{F})t))^2-tr(((h\mathcal{P}+s\mathcal{F})t)^2)-2tr((\mathcal{F}t+\mathcal{N}s)h(f\mathcal{P}+t\mathcal{F})t)\Big]\\
&+6\frac{c_1-c_2}{16}\Big[-tr((h\mathcal{P}+s\mathcal{F})t\mathcal{F}t)-tr((f\mathcal{P}+t\mathcal{F})t\mathcal{N}h)-tr(h\mathcal{P}t(\mathcal{F}t+\mathcal{N}s))\Big]\\
&-3\frac{c_1+c_2}{16}\Big[-2tr((f\mathcal{P}+t\mathcal{F})\mathcal{T}(\mathcal{F}t+\mathcal{N}s)\mathcal{F})-2tr(\mathcal{P}(f\mathcal{P}+t\mathcal{F}))tr(\mathcal{N}(\mathcal{F}t+\mathcal{N}s))\\
&+3tr(t(\mathcal{F}t+\mathcal{N}s)h(f\mathcal{P}+t\mathcal{F}))-2tr((\mathcal{P}t+\mathcal{T}s)(\mathcal{F}t+\mathcal{N}s)(h\mathcal{P}+s\mathcal{F})(f\mathcal{P}+t\mathcal{F}))\\
&-2tr((f\mathcal{P}+t\mathcal{F})^2)\cdot tr((\mathcal{F}t+\mathcal{N}s)^2)\Big]\\
&+6\frac{c_1-c_2}{16}\Big[tr(\mathcal{T}(\mathcal{F}t+\mathcal{N}s)(h\mathcal{P}+s\mathcal{F})(f\mathcal{P}+t\mathcal{F}))\\
&+tr((\mathcal{P}t+\mathcal{T}s)(\mathcal{F}t+\mathcal{N}s)\mathcal{F}(f\mathcal{P}+t\mathcal{F}))\\
&+tr((f\mathcal{P}+t\mathcal{F})^2)\cdot tr(\mathcal{N}(\mathcal{F}t+\mathcal{N}s))+tr((\mathcal{F}t+\mathcal{N}s)^2)\cdot tr(\mathcal{P}(f\mathcal{P}+t\mathcal{F}))\Big]\\
&+\frac{3}{2}\sum_{\alpha}|[f\mathcal{P}+t\mathcal{F},A_\alpha]|^2+3\sum_{\alpha,\beta}g(A_\alpha tv_\beta,A_\beta tv_\alpha)\\
&+\sum_{\alpha}\Big[2tr(A_\alpha^2)-2tr(A_{sv_\alpha}^2)+tr(A_\alpha A_{sv_\alpha}f)-tr(f)tr(A_\alpha A_{sv_\alpha})+tr(f)tr(A_\alpha^2 f)\\
&-2tr(A_\alpha^2 f^2)-2(tr(A_\alpha f))^2+tr((A_\alpha f)^2)+3tr(A_\alpha A_{(h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s)v_\alpha})\Big]\Big\}\\
&+\frac{c_1-c_2}{16}\Big\{l\cdot tr(A_\alpha^2 f)-l\cdot tr(A_\alpha A_{sv_\alpha})\\
&+6\frac{c_1+c_2}{16}\Big[tr(\mathcal{F}\mathcal{T})\cdot tr(\mathcal{F}(\mathcal{P}t+\mathcal{T}s))-tr(\mathcal{F}\mathcal{T}\mathcal{F}(\mathcal{P}t+\mathcal{T}s))\\
&+2tr(\mathcal{N}\mathcal{F}\mathcal{P}(\mathcal{P}t+\mathcal{T}s))+2tr(h\mathcal{T})\cdot tr(h(\mathcal{P}t+\mathcal{T}s))\\
&-tr(h\mathcal{T}h(\mathcal{P}t+\mathcal{T}s))+tr((h\mathcal{P}+s\mathcal{F})\mathcal{T})\cdot tr((h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s))\\
&-tr((h\mathcal{P}+s\mathcal{F})\mathcal{T}(h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s))\\
&+2tr((\mathcal{F}t+\mathcal{N}s)((h\mathcal{P}+s\mathcal{F})(f\mathcal{P}+t\mathcal{F})\mathcal{T})\Big]\\
&+6\frac{c_1-c_2}{16}\Big[(tr(\mathcal{F}(\mathcal{P}t+\mathcal{T}s)))^2-tr(\mathcal{T}\mathcal{F}(\mathcal{P}t+\mathcal{T}s)(h\mathcal{P}+s\mathcal{F}))\\
&-tr(\mathcal{F}\mathcal{T}(h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s))+tr(\mathcal{F}\mathcal{T})\cdot tr((h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s))\\
&+2tr(\mathcal{T}\mathcal{N}(h\mathcal{P}+s\mathcal{F})(f\mathcal{P}+t\mathcal{F}))+2tr((\mathcal{F}t+\mathcal{N}s)(h\mathcal{P}+s\mathcal{F})\mathcal{P}\mathcal{T})\Big]\\
&-3\frac{c_1+c_2}{16}\Big[-2tr(\mathcal{T}\mathcal{N}\mathcal{F}(f\mathcal{P}+t\mathcal{F}))-2tr(\mathcal{P}(f\mathcal{P}+t\mathcal{F}))\cdot tr(\mathcal{N}^2)\\
&+3tr(t\mathcal{N}h(f\mathcal{P}+t\mathcal{F}))-2tr((\mathcal{P}t+\mathcal{T}s)\mathcal{N}(h\mathcal{P}+s\mathcal{F})(f\mathcal{P}+t\mathcal{F}))\\
&-2tr((f\mathcal{P}+t\mathcal{F})^2)\cdot tr(\mathcal{N}(\mathcal{F}t+\mathcal{N}s))\Big]\\
&-3\frac{c_1-c_2}{16}\Big[-2tr(\mathcal{T}\mathcal{N}(h\mathcal{P}+s\mathcal{F})(f\mathcal{P}+t\mathcal{F}))-2tr(\mathcal{N}\mathcal{F}(f\mathcal{P}+t\mathcal{F})(\mathcal{P}t+\mathcal{T}s))\\
&-2tr((f\mathcal{P}+t\mathcal{F})^2)\cdot tr(\mathcal{N}^2)-2tr(\mathcal{P}(f\mathcal{P}+t\mathcal{F}))\cdot tr(\mathcal{N}(\mathcal{F}t+\mathcal{N}s))\Big]
\end{aligned}
$$

$$
\begin{aligned}
&-3\frac{c_1+c_2}{16}\Big[-2tr(\mathcal{F}\mathcal{P}\mathcal{T}(\mathcal{F}t+\mathcal{N}s))-2tr(\mathcal{P}^2)tr(\mathcal{N}(\mathcal{F}t+\mathcal{N}s))\\
&+3tr(h\mathcal{P}t(\mathcal{F}t+\mathcal{N}s))-2tr((\mathcal{P}t+\mathcal{T}s)(\mathcal{F}t+\mathcal{N}s)(h\mathcal{P}+s\mathcal{F})\mathcal{P})\\
&-2tr((f\mathcal{P}+t\mathcal{F})\mathcal{P})\cdot tr((\mathcal{F}t+\mathcal{N}s)^2)\Big]\\
&-3\frac{c_1-c_2}{16}\Big[-4tr(\mathcal{P}\mathcal{T}(\mathcal{F}t+\mathcal{N}s)(h\mathcal{P}+s\mathcal{F}))\\
&-2tr((f\mathcal{P}+t\mathcal{F})\mathcal{P})\cdot tr(\mathcal{N}(\mathcal{F}t+\mathcal{N}s))-2tr(\mathcal{P}^2)\cdot tr((\mathcal{F}t+\mathcal{N}s)^2)\Big]\\
&+6tr(A_\alpha A_{(h\mathcal{P}+s\mathcal{F})\mathcal{T}v_\alpha})-6tr(A_\alpha^2(f\mathcal{P}+t\mathcal{F})\mathcal{P})+6tr(A_\alpha\mathcal{P}A_\alpha(f\mathcal{P}+t\mathcal{F}))\Big\}
\end{aligned}
$$

As in [4], we have the following result:

Lemma 3.2 Let M be an l-dimensional minimal submanifold in $\overline{M}^m\times\overline{M}^n$. If U is a parallel section in the normal bundle of M, then

$$
\begin{aligned}
&div(\nabla_{\mathcal{T}U}\mathcal{T}U)\\
=\;&\frac{c_1+c_2}{16}\Big\{(l-1)g(\mathcal{T}U,\mathcal{T}U)+3g(\mathcal{P}\mathcal{T}U,\mathcal{P}\mathcal{T}U)+2tr(f)g(f\mathcal{T}U,\mathcal{T}U)\\
&-g(f\mathcal{T}U,f\mathcal{T}U)+3g((f\mathcal{P}+t\mathcal{F})\mathcal{T}U,(f\mathcal{P}+t\mathcal{F})\mathcal{T}U)\Big\}\\
&+\frac{c_1-c_2}{16}\Big\{(l-2)g(f\mathcal{T}U,\mathcal{T}U)+tr(f)\cdot g(\mathcal{T}U,\mathcal{T}U)\\
&+6g(\mathcal{P}\mathcal{T}U,(f\mathcal{P}+t\mathcal{F})\mathcal{T}U)\Big\}\\
&+\frac{1}{2}|[\mathcal{P},A_U]|^2+tr(A_{\mathcal{N}U}^2)-2tr(A_UA_{\mathcal{N}U}\mathcal{P})-tr(A_U^2)\\
&+\sum_\alpha g(A_U\mathcal{T}v_\alpha,A_U\mathcal{T}v_\alpha)-\sum_\alpha g(A_\alpha\mathcal{T}U,A_\alpha\mathcal{T}U)
\end{aligned}
\tag{3.51}
$$

Proof From Gauss equation, we get

$$
\begin{aligned}
&S(\mathcal{T}U,\mathcal{T}U)\\
=\;&\frac{c_1+c_2}{16}\Big\{(l-1)g(\mathcal{T}U,\mathcal{T}U)+3g(\mathcal{P}\mathcal{T}U,\mathcal{P}\mathcal{T}U)+2tr(f)g(f\mathcal{T}U,\mathcal{T}U)\\
&-g(f\mathcal{T}U,f\mathcal{T}U)+3g((f\mathcal{P}+t\mathcal{F})\mathcal{T}U,(f\mathcal{P}+t\mathcal{F})\mathcal{T}U)\Big\}\\
&+\frac{c_1-c_2}{16}\Big\{(l-2)g(f\mathcal{T}U,\mathcal{T}U)+tr(f)\cdot g(\mathcal{T}U,\mathcal{T}U)\\
&+6g(\mathcal{P}\mathcal{T}U,(f\mathcal{P}+t\mathcal{F})\mathcal{T}U)\Big\}-\sum_\alpha g(A_\alpha\mathcal{T}U,A_\alpha\mathcal{T}U)
\end{aligned}
\tag{3.52}
$$

where S is the Ricci tensor of M.

From [4] we know that

$$
\nabla_X(\mathcal{T}U)=-\mathcal{P}A_UX+A_{\mathcal{N}U}X \tag{3.53}
$$

Therefore

$$
\begin{aligned}
div(\mathcal{T}U)&=\sum_i g(e_i,\nabla_{e_i}(\mathcal{T}U))=\sum_i g(e_i,-\mathcal{P}A_Ue_i+A_{\mathcal{N}U}e_i)\\
&=-tr(\mathcal{P}A_U)+tr(A_{\mathcal{N}U})=0,
\end{aligned}
\tag{3.54}
$$

$$
\begin{aligned}
&|\nabla\mathcal{T}U|^2=\sum_i g(\nabla_{e_i}\mathcal{T}U,\nabla_{e_i}\mathcal{T}U) \qquad (3.55)\\
=\;& tr(A_U^2)+tr(A_{\mathcal{N}U}^2)-2tr(A_UA_{\mathcal{N}U}\mathcal{P})-\sum_\alpha g(A_U\mathcal{T}v_\alpha,A_U\mathcal{T}v_\alpha),
\end{aligned}
$$

$$
\begin{aligned}
&|L_{\mathcal{T}U}g|^2=\sum_{i,j}\Big[L_{\mathcal{T}U}g(e_i,e_j)\Big]^2=\sum_{i,j}\Big[g(\nabla_{e_i}\mathcal{T}U,e_j)+g(\nabla_{e_j}\mathcal{T}U,e_i)\Big]^2\\
=\;& |[\mathcal{P},A_U]|^2+4tr(A_{\mathcal{N}U}^2)-8tr(A_UA_{\mathcal{N}U}\mathcal{P}) \qquad (3.56)
\end{aligned}
$$

Put (3.26)-(3.30) into

$$
\begin{aligned}
div(\nabla_{\mathcal{T}U}\mathcal{T}U) =\;& div(div(\mathcal{T}U)\mathcal{T}U)+S(\mathcal{T}U,\mathcal{T}U)\\
&+\frac{1}{2}|L_{\mathcal{T}U}g|^2-|\nabla\mathcal{T}U|^2-(div(\mathcal{T}U))^2
\end{aligned}
$$

we get the result. □

Since the normal connection of M is flat, we can choose an orthonormal basis v_α of $T(M)^\perp$ such that $Dv_\alpha=0$ for all α. Thus

$$
\begin{aligned}
&div(\nabla_{\mathcal{T}v_\alpha}\mathcal{T}v_\alpha)\\
=\;& \frac{c_1+c_2}{16}\Big\{(l-1)g(\mathcal{T}v_\alpha,\mathcal{T}v_\alpha)+3g(\mathcal{P}\mathcal{T}v_\alpha,\mathcal{P}\mathcal{T}v_\alpha)+2tr(f)g(f\mathcal{T}v_\alpha,\mathcal{T}v_\alpha)\\
&-g(f\mathcal{T}v_\alpha,f\mathcal{T}v_\alpha)+3g((f\mathcal{P}+t\mathcal{F})\mathcal{T}v_\alpha,(f\mathcal{P}+t\mathcal{F})\mathcal{T}v_\alpha)\Big\} \qquad (3.57)\\
&+\frac{c_1-c_2}{16}\Big\{(l-2)g(f\mathcal{T}v_\alpha,\mathcal{T}v_\alpha)+tr(f)\cdot g(\mathcal{T}v_\alpha,\mathcal{T}v_\alpha)\\
&+6g(\mathcal{P}\mathcal{T}v_\alpha,(f\mathcal{P}+t\mathcal{F})\mathcal{T}v_\alpha)\Big\}\\
&+\frac{1}{2}|[\mathcal{P},A_\alpha]|^2+tr(A_{\mathcal{N}v_\alpha}^2)-2tr(A_\alpha A_{\mathcal{N}v_\alpha}\mathcal{P})-tr(A_\alpha^2)\\
=\;& \frac{c_1+c_2}{16}\Big\{-(l-1)\cdot tr(\mathcal{F}\mathcal{T})+3tr(\mathcal{P}^2\mathcal{T}\mathcal{F})-2tr(f)\cdot tr(f\mathcal{T}\mathcal{F})+tr(f^2\mathcal{T}\mathcal{F})\\
&+3tr((f\mathcal{P}+t\mathcal{F})^2\mathcal{T}\mathcal{F})\Big\}\\
&+\frac{c_1-c_2}{16}\Big\{-(l-2)tr(f\mathcal{T}\mathcal{F})-tr(f)\cdot tr(\mathcal{F}\mathcal{T})+6tr((f\mathcal{P}+t\mathcal{F})\mathcal{T}\mathcal{F}\mathcal{P})\Big\}\\
&+\frac{1}{2}|[\mathcal{P},A_\alpha]|^2+tr(A_{\mathcal{N}v_\alpha}^2)-2tr(A_\alpha A_{\mathcal{N}v_\alpha}\mathcal{P})-tr(A_\alpha^2)
\end{aligned}
$$

Lemma 3.3 Let M be an l-dimensional minimal submanifold in $\overline{M}^m\times\overline{M}^n$. If U is a parallel section in the normal bundle of M, then

$$
\begin{aligned}
&div(\nabla_{tU}tU) \qquad (3.58)\\
=\;& -div(tr(A_Uf)tU)+\frac{c_1+c_2}{16}\Big\{(l-1)g(tU,tU)+3g(\mathcal{P}tU,\mathcal{P}tU)\\
&+2tr(f)g(ftU,tU)-g(ftU,ftU)+3g((f\mathcal{P}+t\mathcal{F})tU,(f\mathcal{P}+t\mathcal{F})tU)\Big\}\\
&+\frac{c_1-c_2}{16}\Big\{(l-2)g(ftU,tU)+tr(f)\cdot g(tU,tU)+6g(\mathcal{P}tU,(f\mathcal{P}+t\mathcal{F})tU)\Big\}\\
&+tr(A_{sU}^2)-tr(A_U^2)+tr(A_U^2f^2)+tr((A_Uf)^2)-(tr(A_Uf))^2\\
&-2tr(A_UA_{sU}f)+\sum_\alpha g(A_Utv_\alpha,A_Utv_\alpha)-\sum_\alpha g(A_\alpha tU,A_\alpha tU)
\end{aligned}
$$

Proof As in the proof of Lemma 3.2, we have

$$\begin{aligned}
& S(tU,tU) \\
= \ & \frac{c_1+c_2}{16}\Big\{(l-1)g(tU,tU)+3g(\mathcal{P}tU,\mathcal{P}tU)+2tr(f)g(ftU,tU) \\
& -g(ftU,ftU)+3g((f\mathcal{P}+t\mathcal{F})tU,(f\mathcal{P}+t\mathcal{F})tU)\Big\} \\
& +\frac{c_1-c_2}{16}\Big\{(l-2)g(ftU,tU)+tr(f)\cdot g(tU,tU)+6g(\mathcal{P}tU,(f\mathcal{P}+t\mathcal{F})tU)\Big\} \\
& -\sum_\alpha g(A_\alpha tU,A_\alpha tU). \qquad (3.59)
\end{aligned}$$

$$\nabla_X(tU)=A_{sU}X-fA_UX \qquad (3.60)$$

$$div(tU)=-tr(A_Uf) \qquad (3.61)$$

$$|\nabla tU|^2=tr(A_{sU}^2)+tr(A_U^2)-2tr(A_UA_{sU}f)-\sum_\alpha g(A_Utv_\alpha,A_Utv_\alpha) \qquad (3.62)$$

$$|L_{tU}g|^2=4tr(A_{sU}^2)+2tr(A_U^2f^2)+2tr((A_Uf)^2)-8tr(A_UA_{sU}f) \qquad (3.63)$$

Put (3.31), (3.33)-(3.35) into

$$\begin{aligned}
div(\nabla_{tU}tU) \ = \ & div(div(tU)tU)+S(tU,tU) \\
& +\frac{1}{2}|L_{tU}g|^2-|\nabla tU|^2-(div(tU))^2
\end{aligned}$$

we get the result. □

As before we choose an orthonormal basis v_α of $T(M)^\perp$ such that $Dv_\alpha=0$ for all α. Thus

$$\begin{aligned}
& div(\nabla_{tv_\alpha}tv_\alpha) \qquad (3.64) \\
= \ & -div(tr(A_\alpha f)tv_\alpha)+\frac{c_1+c_2}{16}\Big\{(l-1)g(tv_\alpha,tv_\alpha)+3g(\mathcal{P}tv_\alpha,\mathcal{P}tv_\alpha) \\
& +2tr(f)g(ftv_\alpha,tv_\alpha)-g(ftv_\alpha,ftv_\alpha)+3g((f\mathcal{P}+t\mathcal{F})tv_\alpha,(f\mathcal{P}+t\mathcal{F})tv_\alpha)\Big\} \\
& +\frac{c_1-c_2}{16}\Big\{(l-2)g(ftv_\alpha,tv_\alpha)+tr(f)\cdot g(tv_\alpha,tv_\alpha)+6g(\mathcal{P}tv_\alpha,(f\mathcal{P}+t\mathcal{F})tv_\alpha)\Big\} \\
& +tr(A_{sv_\alpha}^2)-tr(A_\alpha^2)+tr(A_\alpha^2f^2)+tr((A_\alpha f)^2)-(tr(A_\alpha f))^2-2tr(A_{v_\alpha}A_{sv_\alpha}f) \\
= \ & -div(tr(A_\alpha f)tv_\alpha)+\frac{c_1+c_2}{16}\Big\{(l-1)\cdot tr(ht)-3tr(\mathcal{P}^2th)+2tr(f)\cdot tr(hft) \\
& -tr(f^2th)-3tr((f\mathcal{P}+t\mathcal{F})^2th)\Big\} \\
& +\frac{c_1-c_2}{16}\Big\{(l-2)\cdot tr(hft)+tr(f)\cdot tr(ht)-6tr((f\mathcal{P}+t\mathcal{F})th\mathcal{P})\Big\} \\
& +tr(A_{sv_\alpha}^2)-tr(A_\alpha^2)+tr(A_\alpha^2f^2)+tr((A_\alpha f)^2)-(tr(A_\alpha f))^2-2tr(A_{v_\alpha}A_{sv_\alpha}f)
\end{aligned}$$

Theorem 3.4 If the submanifold M is compact minimal submanifold in $\overline{M}^m \times \overline{M}^n$, then

$$\begin{aligned}
&\int_M |\nabla A|^2 d\mu + 3(\frac{c_1+c_2}{16})^2 \int_M W_1 d\mu + 6(\frac{c_1-c_2}{16})^2 \int_M W_2 d\mu \\
= & -\int_M \sum_{i,j,\alpha} g((R(e_i,e_j)A_\alpha)e_i, A_\alpha e_j) d\mu \qquad (3.65)\\
&+3(\frac{c_1+c_2}{16})^2 \int_M W_3 d\mu + 6(\frac{c_1-c_2}{16})^2 \int_M W_4 d\mu + \frac{c_1^2-c_2^2}{16^2} \int_M W_5 d\mu \\
&+\frac{c_1+c_2}{16} \int_M W_6 d\mu + \frac{c_1-c_2}{16} \int_M W_7 d\mu
\end{aligned}$$

where

$$\begin{aligned}
W_1 = & -tr(f^2\mathcal{T}\mathcal{F}) + [tr(\mathcal{F}(\mathcal{P}t+\mathcal{T}s))]^2 + 2[tr(h(\mathcal{P}t+\mathcal{T}s))]^2 \qquad (3.66)\\
&+[tr((h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s))]^2 d\mu + [tr(\mathcal{F}\mathcal{T})]^2 + [tr(h\mathcal{P}+s\mathcal{F})\mathcal{T}]^2 \\
&+2[tr(ht)]^2 + [tr((h\mathcal{P}+s\mathcal{F})t)]^2 + 2tr((f\mathcal{P}+t\mathcal{F})^2)\cdot tr((\mathcal{F}t+\mathcal{N}s)^2)
\end{aligned}$$

$$\begin{aligned}
W_2 = & [tr(\mathcal{F}(\mathcal{P}t+\mathcal{T}s))]^2 + tr(\mathcal{F}\mathcal{T})\cdot tr((h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s)) \\
&+tr((f\mathcal{P}+t\mathcal{F})^2)\cdot tr(\mathcal{N}^2) + tr(\mathcal{P}^2)\cdot tr((\mathcal{F}t+\mathcal{N}s)^2) \qquad (3.67)
\end{aligned}$$

$$\begin{aligned}
W_3 = & -(l-1)\cdot tr(\mathcal{F}\mathcal{T}) + 3tr(\mathcal{P}^2\mathcal{T}\mathcal{F}) - 2tr(f)\cdot tr(f\mathcal{T}\mathcal{F}) \\
&+3tr((f\mathcal{P}+t\mathcal{F})^2\mathcal{T}\mathcal{F}) + tr[(\mathcal{F}(\mathcal{P}t+\mathcal{T}s))^2] \\
&-2tr(\mathcal{N}(h\mathcal{P}+s\mathcal{F})\mathcal{P}(\mathcal{P}t+\mathcal{T}s)) + tr[(h(\mathcal{P}t+\mathcal{T}s))^2] \\
&+tr[((h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s))^2] \qquad (3.68)\\
&-2tr[(\mathcal{F}t+\mathcal{N}s)(h\mathcal{P}+s\mathcal{F})(f\mathcal{P}+t\mathcal{F})(\mathcal{P}t+\mathcal{T}s)] \\
&+tr[(\mathcal{F}\mathcal{T})^2] - 2tr(\mathcal{N}\mathcal{F}\mathcal{P}\mathcal{T}) + tr[(\mathcal{F}t)^2] \\
&+tr[(h\mathcal{P}+s\mathcal{F})\mathcal{T}]^2 - 2tr[(\mathcal{F}t+\mathcal{N}s)\mathcal{F}(f\mathcal{P}+t\mathcal{F})\mathcal{T}] \\
&+tr((\mathcal{F}t)^2) + 2tr(\mathcal{N}h\mathcal{P}t) + tr((ht)^2) \\
&+tr(((h\mathcal{P}+s\mathcal{F})t)^2) + 2tr((\mathcal{F}t+\mathcal{N}s)h(f\mathcal{P}+t\mathcal{F})t) \\
&-2tr((f\mathcal{P}+t\mathcal{F})\mathcal{T}(\mathcal{F}t+\mathcal{N}s)\mathcal{F}) \\
&-2tr(\mathcal{P}(f\mathcal{P}+t\mathcal{F}))tr(\mathcal{N}(\mathcal{F}t+\mathcal{N}s)) + 3tr(t(\mathcal{F}t+\mathcal{N}s)h(f\mathcal{P}+t\mathcal{F})) \\
&-2tr((\mathcal{P}t+\mathcal{T}s)(\mathcal{F}t+\mathcal{N}s)(h\mathcal{P}+s\mathcal{F})(f\mathcal{P}+t\mathcal{F}))
\end{aligned}$$

$$\begin{aligned}
W_4 = & -\Big[-tr(\mathcal{T}\mathcal{F}(\mathcal{P}t+\mathcal{T}s)(h\mathcal{P}+s\mathcal{F})) - tr(\mathcal{F}\mathcal{T}(h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s)) \\
&+2tr(\mathcal{T}\mathcal{N}(h\mathcal{P}+s\mathcal{F})(f\mathcal{P}+t\mathcal{F})) + 2tr((\mathcal{F}t+\mathcal{N}s)(h\mathcal{P}+s\mathcal{F})\mathcal{P}\mathcal{T})\Big] \\
&-\Big[tr(\mathcal{T}\mathcal{N}(h\mathcal{P}+s\mathcal{F})(f\mathcal{P}+t\mathcal{F})) + tr(\mathcal{N}\mathcal{F}(f\mathcal{P}+t\mathcal{F})(\mathcal{P}t+\mathcal{T}s)) \\
&+tr(\mathcal{P}(f\mathcal{P}+t\mathcal{F}))\cdot tr(\mathcal{N}(\mathcal{F}t+\mathcal{N}s))\Big] \\
&-\Big[2tr(\mathcal{P}\mathcal{T}(\mathcal{F}t+\mathcal{N}s)(h\mathcal{P}+s\mathcal{F})) \qquad (3.69)\\
&+tr((f\mathcal{P}+t\mathcal{F})\mathcal{P})\cdot tr(\mathcal{N}(\mathcal{F}t+\mathcal{N}s))\Big]
\end{aligned}$$

$$
\begin{aligned}
W_6 &= -\frac{3}{2}\sum_{\alpha}|[f\mathcal{P}+t\mathcal{F},A_\alpha]|^2-3\sum_{\alpha,\beta}g(A_\alpha tv_\beta,A_\beta tv_\alpha)\\
&\quad-\sum_{\alpha}\Big[2tr(A_\alpha^2)-2tr(A_{sv_\alpha}^2)+tr(A_\alpha A_{sv_\alpha}f)-tr(f)tr(A_\alpha A_{sv_\alpha})\\
&\quad+tr(f)tr(A_\alpha^2 f)-2tr(A_\alpha^2 f^2)-2(tr(A_\alpha f))^2+tr((A_\alpha f)^2)\\
&\quad+3tr(A_\alpha A_{(h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s)v_\alpha})\Big] \qquad (3.70)
\end{aligned}
$$

$$
\begin{aligned}
W_5 &= 3\Big[-(l-2)tr(f\mathcal{T}\mathcal{F})-tr(f)\cdot tr(\mathcal{F}\mathcal{T})+6tr((f\mathcal{P}+t\mathcal{F})\mathcal{T}\mathcal{F}\mathcal{P})\Big]\\
&\quad-6\Big[tr(\mathcal{F}(\mathcal{P}t+\mathcal{T}s))\cdot tr((h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s))\\
&\quad-tr(\mathcal{F}(\mathcal{P}t+\mathcal{T}s)(h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s))\\
&\quad+tr(\mathcal{N}(h\mathcal{P}+s\mathcal{F})(f\mathcal{P}+t\mathcal{F})(\mathcal{P}t+\mathcal{T}s)) \qquad (3.71)\\
&\quad+tr((\mathcal{F}t+\mathcal{N}s)(h\mathcal{P}+s\mathcal{F})\mathcal{P}(\mathcal{P}t+\mathcal{T}s))\Big]\\
&\quad-6\Big[tr(\mathcal{F}\mathcal{T})\cdot tr((h\mathcal{P}+s\mathcal{F})\mathcal{T})-tr(\mathcal{F}\mathcal{T}(h\mathcal{P}+s\mathcal{F})\mathcal{T})\\
&\quad+tr(\mathcal{N}\mathcal{F}(f\mathcal{P}+t\mathcal{F})\mathcal{T})+tr(\mathcal{F}\mathcal{P}\mathcal{T}(\mathcal{F}t+\mathcal{N}s))\Big]\\
&\quad-6\Big[-tr((h\mathcal{P}+s\mathcal{F})t\mathcal{F}t)-tr((f\mathcal{P}+t\mathcal{F})t\mathcal{N}h)-tr(h\mathcal{P}t(\mathcal{F}t+\mathcal{N}s))\Big]\\
&\quad-6\Big[tr(\mathcal{T}(\mathcal{F}t+\mathcal{N}s)(h\mathcal{P}+s\mathcal{F})(f\mathcal{P}+t\mathcal{F}))\\
&\quad+tr((\mathcal{P}t+\mathcal{T}s)(\mathcal{F}t+\mathcal{N}s)\mathcal{F}(f\mathcal{P}+t\mathcal{F}))\\
&\quad+tr((f\mathcal{P}+t\mathcal{F})^2)\cdot tr(\mathcal{N}(\mathcal{F}t+\mathcal{N}s))\\
&\quad+tr((\mathcal{F}t+\mathcal{N}s)^2)\cdot tr(\mathcal{P}(f\mathcal{P}+t\mathcal{F}))\Big]\\
&\quad-6\Big[tr(\mathcal{F}\mathcal{T})\cdot tr(\mathcal{F}(\mathcal{P}t+\mathcal{T}s))-tr(\mathcal{F}\mathcal{T}\mathcal{F}(\mathcal{P}t+\mathcal{T}s))\\
&\quad+2tr(\mathcal{N}\mathcal{F}\mathcal{P}(\mathcal{P}t+\mathcal{T}s))+2tr(h\mathcal{T})\cdot tr(h(\mathcal{P}t+\mathcal{T}s))\\
&\quad-tr(h\mathcal{T}h(\mathcal{P}t+\mathcal{T}s))+tr((h\mathcal{P}+s\mathcal{F})\mathcal{T})\cdot tr((h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s))\\
&\quad-tr((h\mathcal{P}+s\mathcal{F})\mathcal{T}(h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s))\\
&\quad+2tr((\mathcal{F}t+\mathcal{N}s)(h\mathcal{P}+s\mathcal{F})(f\mathcal{P}+t\mathcal{F})\mathcal{T})\Big]\\
&\quad+3\Big[-2tr(\mathcal{T}\mathcal{N}\mathcal{F}(f\mathcal{P}+t\mathcal{F}))-2tr(\mathcal{P}(f\mathcal{P}+t\mathcal{F}))\cdot tr(\mathcal{N}^2)\\
&\quad+3tr(t\mathcal{N}h(f\mathcal{P}+t\mathcal{F}))-2tr((\mathcal{P}t+\mathcal{T}s)\mathcal{N}(h\mathcal{P}+s\mathcal{F})(f\mathcal{P}+t\mathcal{F}))\\
&\quad-2tr((f\mathcal{P}+t\mathcal{F})^2)\cdot tr(\mathcal{N}(\mathcal{F}t+\mathcal{N}s))\Big]\\
&\quad+3\Big[-2tr(\mathcal{F}\mathcal{P}\mathcal{T}(\mathcal{F}t+\mathcal{N}s))-2tr(\mathcal{P}^2)tr(\mathcal{N}(\mathcal{F}t+\mathcal{N}s))\\
&\quad+3tr(h\mathcal{P}t(\mathcal{F}t+\mathcal{N}s))-2tr((\mathcal{P}t+\mathcal{T}s)(\mathcal{F}t+\mathcal{N}s)(h\mathcal{P}+s\mathcal{F})\mathcal{P})\\
&\quad-2tr((f\mathcal{P}+t\mathcal{F})\mathcal{P})\cdot tr((\mathcal{F}t+\mathcal{N}s)^2)\Big]
\end{aligned}
$$

$$
\begin{aligned}
W_7 &= -l\cdot\Big[tr(A_\alpha^2 f)-tr(A_\alpha A_{sv_\alpha})\Big]-6\Big[tr(A_\alpha A_{(h\mathcal{P}+s\mathcal{F})\mathcal{T}v_\alpha})\\
&\quad-tr(A_\alpha^2(f\mathcal{P}+t\mathcal{F})\mathcal{P})+tr(A_\alpha\mathcal{P}A_\alpha(f\mathcal{P}+t\mathcal{F}))\Big] \qquad (3.72)
\end{aligned}
$$

Note that all the terms in W_1 and W_2 are positive, the left side of (3.39) is positive, therefore we get the following

Corollary 3.5 Let M be an l-dimensional compact minimal submanifold in $\overline{M}^m \times \overline{M}^n$ with flat normal connection. If the sectional curvature K of M has a lower bound C and if the second fundamental form A satisfies

$$\begin{aligned} &-l\cdot C\sum_\alpha tr(A_\alpha^2)+3(\frac{c_1+c_2}{16})^2W_3+6(\frac{c_1-c_2}{16})^2W_4\\ &+\frac{c_1^2-c_2^2}{16^2}W_5+\frac{c_1+c_2}{16}W_6+\frac{c_1-c_2}{16}W_7<0 \end{aligned} \tag{3.73}$$

then M is totally geodesic.

Proof. We can choosing an orthonormal basis $\{e_i\}$ of T_xM such that $A_\alpha e_i = h_i^\alpha e_i$, $i=1,2,\cdots$, then

$$\begin{aligned} &\sum_{i,j}((R(e_i,e_j)A)_\alpha e_i, A_\alpha e_j) \qquad (3.74)\\ =\ &\sum_{i,j} g(R(e_i,e_j)A_\alpha e_i, A_\alpha e_j)-\sum_{i,j} g(A_\alpha R(e_i,e_j)e_i, A_\alpha e_j)\\ =\ &\frac{1}{2}\sum_{i,j}(h_i^\alpha-h_j^\alpha)^2K_{ij}\\ \geq\ &\frac{C}{2}\sum_{i,j}(h_i^\alpha-h_j^\alpha)^2=l\cdot C\cdot tr(A_\alpha^2) \end{aligned}$$

where K_{ij} denotes the sectional curvature of M with respect to the section spanned by e_i and e_j.

Denote the left side of (3.39) by $\mathcal{L}$, then there exist a small $\varepsilon>0$, such that

$$\begin{aligned} &\mathcal{L}+\varepsilon\int_M\sum_\alpha tr(A_\alpha^2)d\mu\\ &\leq\int_M\Big[(\varepsilon-l\cdot C)\sum_\alpha tr(A_\alpha^2)+3(\frac{c_1+c_2}{16})^2W_3+6(\frac{c_1-c_2}{16})^2W_4\\ &+\frac{c_1^2-c_2^2}{16^2}W_5+\frac{c_1+c_2}{16}W_6+\frac{c_1-c_2}{16}W_7\Big]d\mu\\ &\leq 0 \end{aligned} \tag{3.75}$$

Therefore $\sum_\alpha tr(A_\alpha^2)=0$. □

Corollary 3.6 Let M be an l-dimensional compact minimal submanifold in $\overline{M}^m\times\overline{M}^n$ with flat normal connection, $c_1+c_2>0$. If the sectional curvature K of M has a lower bound C and if the second fundamental form A satisfies

$$\begin{aligned} &-l\cdot C\sum_\alpha tr(A_\alpha^2)+3(\frac{c_1+c_2}{16})^2W_3+6(\frac{c_1-c_2}{16})^2W_4 \qquad (3.76)\\ &+\frac{c_1^2-c_2^2}{16^2}W_5+\frac{c_1+c_2}{16}W_6'+\frac{c_1-c_2}{16}W_7\leq 0, \end{aligned}$$

where $W_6'=W_6+2\sum_\alpha tr(A_\alpha^2)+\frac{3}{2}\sum_\alpha|[f\mathcal{P}+t\mathcal{F},A_\alpha]|^2$, then M is totally geodesic.

Proof. We only need to move the terms $-2\sum_\alpha tr(A_\alpha^2)$ and $-\frac{3}{2}\sum_\alpha |[f\mathcal{P}+t\mathcal{F}, A_\alpha]|^2$ on the left side of (3.39). The rest are the same as the proof of Corollary 3.5. □

4 The F-anti invariant submanifold

From now on we assume that M is F-anti invariant, that is $FT_xM \subset T_x(M)^\perp$, or equivalently $f \equiv 0$. Then from Lemma 3.2 or (3.31) we have

$$\begin{aligned}
&\sum_\alpha \Big[\frac{1}{2}|[\mathcal{P}, A_\alpha]|^2 + tr(A^2_{\mathcal{N}v_\alpha}) - 2tr(A_\alpha A_{\mathcal{N}v_\alpha}\mathcal{P}) - tr(A_\alpha^2)\Big] \\
= \;& -\frac{c_1+c_2}{16}\sum_\alpha \Big[(l-1)g(\mathcal{T}v_\alpha, \mathcal{T}v_\alpha) + 3g(\mathcal{P}\mathcal{T}v_\alpha, \mathcal{P}\mathcal{T}v_\alpha) + 3g(t\mathcal{F}\mathcal{T}v_\alpha, t\mathcal{F}\mathcal{T}v_\alpha)\Big] \\
& -6\frac{c_1-c_2}{16}\sum_\alpha g(\mathcal{P}\mathcal{T}v_\alpha, t\mathcal{F}\mathcal{T}v_\alpha) + \sum_\alpha div(\nabla_{\mathcal{T}v_\alpha}\mathcal{T}v_\alpha) \qquad (4.77) \\
= \;& -\frac{c_1+c_2}{16}\Big[-(l-1)tr(\mathcal{F}\mathcal{T}) + 3tr(\mathcal{P}^2\mathcal{T}\mathcal{F}) + 3tr(ht(\mathcal{F}\mathcal{T})^2)\Big] \\
& -6\frac{c_1-c_2}{16}tr(t\mathcal{F}\mathcal{T}\mathcal{F}\mathcal{P}) + \sum_\alpha div(\nabla_{\mathcal{T}v_\alpha}\mathcal{T}v_\alpha)
\end{aligned}$$

and from Lemma 3.3 or (3.38) we have

$$\begin{aligned}
&\sum_\alpha \Big[tr(A_\alpha^2) - tr(A^2_{sv_\alpha})\Big] \\
= \;& \frac{c_1+c_2}{16}\sum_\alpha \Big[(l-1)g(tv_\alpha, tv_\alpha) + 3g(\mathcal{P}tv_\alpha, \mathcal{P}tv_\alpha) + 3g(t\mathcal{F}tv_\alpha, t\mathcal{F}tv_\alpha)\Big] \\
& +6\frac{c_1-c_2}{16}\sum_\alpha g(\mathcal{P}tv_\alpha, t\mathcal{F}tv_\alpha) - \sum_\alpha div(\nabla_{tv_\alpha}tv_\alpha) \qquad (4.78) \\
= \;& \frac{c_1+c_2}{16}\Big[(l-1)tr(ht) - 3tr(\mathcal{P}^2th) - 3tr(h\mathcal{T}ht\mathcal{F}t)\Big] \\
& -6\frac{c_1-c_2}{16}tr(t\mathcal{F}th\mathcal{P}) - \sum_\alpha div(\nabla_{tv_\alpha}tv_\alpha)
\end{aligned}$$

Theorem 4.1 Let M be an F-anti invariant submanifold in an arbitrary kaehler product manifold $\overline{M}^m \times \overline{M}^n$, if $s \equiv 0$, then M is totally geodesic.

Proof. From [7] Theorem 3.3 on Page 426, we know that when M is F-anti invariant, $A_{hX}Y = 0$ or equivalently $tB(X,Y) = 0$ for all tangent vector fields X, Y of M. Then by (3.14)

$$\sum_\alpha \Big[tr(A_\alpha^2) - tr(A^2_{sv_\alpha})\Big] = 0.$$

Therefore $\sum_\alpha tr(A_\alpha^2) = \sum_\alpha tr(A^2_{sv_\alpha}) = 0$. □

Similarly

$$\sum_{\alpha,\beta} g(A_\alpha tv_\beta, A_\beta tv_\alpha) = \sum_{i,\alpha} g(e_i, A_{hA_\alpha e_i}tv_\alpha) = 0.$$

Theorem 4.2 Let M be an F-anti invariant Lagrangian submanifold in an arbitrary kaehler product manifold $\overline{M}^m \times \overline{M}^n$, then M is totally geodesic.

Proof. The dimension of Lagrangian submanifold is half of that of $\overline{M}^m \times \overline{M}^n$, by Theorem 3.3 on Page 426 of [7], we get the result. □

Remark 4.3 Certainly Corollary 3.5, Corollary 3.6 are still true in this case and $W_1, \cdots, W_7$ have simpler expressions, specially

$$W_6 = -\frac{3}{2}\sum_{\alpha} |[t\mathcal{F}, A_\alpha]|^2 - 3\sum_{\alpha} tr(A_\alpha A_{(h\mathcal{P}+s\mathcal{F})(\mathcal{P}t+\mathcal{T}s)v_\alpha})$$

References

[1] M.Atceken, F-invariant submanifolds of kaehlerian product manifold, *Turk.J.Math.*,**28** (2004),367-381.

[2] M.Atceken and S.Keles, Two theorems on invariant submanifolds of product Riemannian manifold.*Indian J.Pure and Appl.Math.*,**34**7(2003),no.7,1035-1044.

[3] M.A.Choudhary, Totally real submanifolds of Kaehler product manifold. *J.Egypt. Math.Soc.*,**24** (2016),258-262.

[4] M.Kon, Pinching theorems for a compact minimal submanifold in a complex projective space. *Bull.Austral.Math.Soc.*,**77**(2008),99-114.

[5] J.Simons, Minimal varieties in Riemannian manifolds, *Ann.of Math.*,**88**(1968),62-105.

[6] Xu Senlin and Ni Yilong, Submanifolds of product Riemannian manifold, *Acta Mathematica Scientia*,**20 B** (2000),213-218.

[7] K.Yano and M.Kon, *Structures on manifolds*,World Scientific, Singapore, 1984.